

The Skin Cancer Surgery Center

Skin Cancer

Skin Cancer is by far the most common malignant tumor in humans. The most common types of skin cancer are *basal cell carcinoma, squamous cell carcinoma, and melanoma*. These begin as a single point in the upper layers of the skin and slowly enlarge, spreading both along the surface and downward. These extensions cannot always be directly seen. The tumor often extends far beyond what is visible on the surface of the skin. Skin cancer may invade and destroy structures in their path. Although these skin cancers are locally destructive, basal cell and squamous cell carcinomas do not tend to metastasize (spread) to distant parts of the body. Metastasis of basal cell carcinoma is extremely rare and usually occurs only in the setting of long-standing large tumors where the patient's immune system is compromised. Squamous cell carcinoma is slightly more dangerous, and patients must be observed for any spread of the tumor. Such spread is still infrequent. Melanoma is a very different and more dangerous kind of skin cancer. Other testing and close long-term follow-up is warranted after its removal.

Excessive exposure to sunlight is the single most important factor associated with the development of skin cancers. In addition, the tendency to develop these cancers appears to be hereditary in certain ethnic groups, especially those with fair complexions and poor tanning abilities. Fair-skinned people develop skin cancers more frequently than dark-skinned people, and the more sun exposure they receive, the more likely they are to develop a skin cancer. Other factors, including exposure to radiation, trauma and exposure to certain chemicals, may also be involved in the development of skin cancers.

The vast majority of skin cancers are present for more than a year before being diagnosed and their growth is rather slow. Skin cancers may be more aggressive in certain instances: patients whose immune system is compromised, patients with a medical history of leukemia or lymphoma, cancers in certain locations such as the ear, lips, lower nose, or around the eyes.

Mohs Micrographic Surgery

There are five standard methods for the treatment of skin cancers. The two nonsurgical treatments are cryotherapy (deep freezing) and radiation therapy. The three surgical methods include simple excision, physical destruction (curettage with electrodesiccation) and Mohs Micrographic Surgery. Newer methods include photodynamic therapy and immunochemotherapy.

In 1935, Dr. Frederic Mohs developed a technique for cancer removal known as chemosurgery. Originally, chemicals were applied to the skin during the surgery. These chemicals are now rarely used, but the name Mohs chemosurgery continues to be associated with the procedure now correctly termed Mohs Micrographic Surgery.

After the removal of the visible portion of the tumor by excision or curettage (debulking), there are two basic steps to each Mohs Micrographic Surgery stage. First, a thin layer of tissue is surgically excised from the base of the site. This layer is generally only 1-2 mm larger than the clinical tumor. Next, this tissue is mapped and processed in a unique manner and examined under the microscope. Our doctors examine the entire bottom surface and outside edges of the tissue on the microscopic slides. (This differs from the "frozen sections" prepared in a hospital setting which, in fact, represent only a tiny sampling of the tumor margins.) This tissue has been marked to orient top to bottom and left to right. If any tumor is seen during the microscopic examination, its location is established, and a thin layer of additional tissue is excised from the involved area. The microscopic examination is then repeated. The entire process is repeated until no tumor is found.

Mohs Micrographic Surgery allows for the selective removal of the skin cancer with the preservation of as much of the surrounding normal tissue as is possible. Because of this complete systematic microscopic search for the "roots" of the skin cancer, Mohs Micrographic Surgery offers *the highest chance for complete removal of the cancer* while sparing the normal tissue. The cure rate for new basal cell and squamous cell carcinomas exceeds 98%. As a result, Mohs Micrographic Surgery is very useful for large tumors, tumors with indistinct borders, tumors near vital functional or cosmetic structures, and tumors for which other forms of therapy have failed. No surgeon or technique can guarantee 100% chance of cure.

Preoperative Visit

If you would feel more comfortable meeting Drs. Welch, Toner, Liu, or Montemarano and our staff, we welcome the opportunity for a preoperative consultation. However, if you feel comfortable with the explanation provided by this pamphlet and your referring doctor, feel free to schedule your surgery directly. If you take *Coumadin* or *blood pressure medications* or if you normally take an *antibiotic prior to dental work*, please bring this to our attention so that there is no delay in your surgery.

Before Mohs Micrographic Surgery


Be well rested and eat a good breakfast, unless instructed otherwise. Take your usual medications, unless directed otherwise. Please do not take Ibuprofen (Motrin, Advil, etc.) or **Vitamin E supplements**. These medications may “thin” your blood and cause more bleeding. You may substitute acetaminophen (Tylenol) if required. Do not drink any alcoholic beverages for 24 hours before surgery.

Shampoo your hair before surgery, as your wound and initial dressing may have to remain dry for 24 hours thereafter. The length of the procedure varies depending on the size and location of the skin cancer and the type of reconstruction to be done. Although the average time is 3-6 hours, **you should plan on spending much of the day in our office**. We ask that you limit the number of people accompanying you to one other because of the limited space in our waiting room. There is plenty of time spent waiting for the lab work, so bring a book or handiwork to keep busy.



The Day of Surgery

Appointments for surgery are scheduled throughout the day. It is a good idea to wear loose fitting clothing and avoid “pullover” clothing. Tylenol may be taken before surgery in order to reduce discomfort. Also, if the operative site is on the face, please do not wear make-up on or around the area and **do not use any fragrance or scented toiletries**. We will obtain your written consent for the procedure, photographs will be taken, and your blood pressure will be recorded. If you have any additional questions, please feel free to ask them at this time.



The area surrounding the skin cancer will be cleansed with an anti-bacterial soap. The doctor will then anesthetize (numb) the area of skin containing the cancer by a small local injection. This injection will probably be similar to the one you received for your biopsy. We will be as gentle as we can when administering this. It usually takes 15 minutes to anesthetize the involved area and remove the tissue. After the tissue has been removed, it will be processed in our office laboratory.





An injection numbs the area. The visible portion of the tumor is debulked.



A thin layer of tissue is excised from the surrounding skin and base. The removed tissue is mapped and sectioned.



The deep and peripheral margins of each section are thinly sliced with a microtome and mounted on microscope slides for examination



If additional tumor is found, it is located on the map, marked and removed. The examination/removal process continues until no tumor is found.

Depending upon the amount of tissue removed, processing usually takes an additional 30-60 minutes. Your wound will be bandaged, and you will move to the waiting room while the tissue is processed, stained, and examined by the doctor. If the microscopic examination of the removed tissue reveals the presence of additional tumor, we will go back and remove more tissue. The Mohs technique allows us to precisely map out where the roots of the cancer remain. Most skin cancers are removed in 1-3 surgical stages.

Reconstruction

After the skin cancer has been completely removed, a decision is made on the best method for treating the wound created by the surgery. These methods include letting the wound heal by itself, closing the wound in a side to side fashion with stitches, closing the wound with a skin graft or a flap. The best method is determined on an individual basis after the final defect is known. Rarely, other surgical specialists may be utilized for their unique skills if warranted. This is the rare exception rather than the rule as nearly all wound repairs are performed immediately.

After Mohs Micrographic Surgery

Your surgical wound will likely require care during the weeks following surgery. Bruising and swelling, for up to a week, is common after surgery. Detailed written instructions will be provided. Plan to return to our office in 5-7 days. You should also plan on wearing a bandage and avoiding strenuous physical activity for a week.

Most of our patients report minimal pain which responds readily to Tylenol. You may experience a sensation of tightness across the area of surgery. Skin cancers frequently involve nerves and months may pass before your skin sensation returns to normal. In rare instances, the numbness may be permanent. You may also experience itching after your wound has healed. Complete healing of the surgical scar takes place over 12-18 months. Especially during the first few months, the site may feel "thick," swollen, or lumpy, and there may be some redness. Gentle massage of the area (starting no sooner than 1 month after the surgery) will speed the healing process.

An indefinite follow-up period of observation is necessary after the wound has healed. You may be asked to return to ensure proper healing.

Studies have shown that once you develop a skin cancer, there is a strong possibility of developing other skin cancers in the future. Should you notice any suspicious areas, it is best to check with your physician for a complete evaluation. You will be reminded to return to your dermatologist on a frequent basis for continued surveillance of your skin.




Risks of Mohs Micrographic Surgery


Because each patient is unique, it is impossible to discuss all the possible complications and risks in this format. The usual risks are discussed below. Our doctors will discuss any additional problems associated with your particular case. *Please understand that these occurrences are the exception and not the rule.*

- The defect created by the removal of the skin cancer may be larger than anticipated. There is no way to predict prior to surgery the exact size of the final defect.
- There will be a scar at the site of the removal. We will make every effort to obtain optimal cosmetic results, but our primary goal is to remove the entire tumor. Again, Mohs surgery will leave you with the *smallest wound* thus creating the best opportunity for optimal cosmetic results.
- There may be poor wound healing. At times, despite our best efforts, for various reasons (such as bleeding, poor physical condition, smoking, diabetes, or other diseases), healing is slow or the wound may reopen. Flaps and grafts utilized to repair the defect may at times fail. Under these circumstances, the wound will usually be left to heal on its own. The scar may be revised at a later date if necessary.
- There may be a loss of motor (muscle) or sensory (feeling) nerve function. **Rarely**, the tumor invades nerve fibers. When this is the case, the nerves must be removed along with the tumor. Prior to your surgery, the doctor will discuss with you any major nerves which might be near your tumor.
- The tumor may involve an important structure. Many are near or on vital structures such as the eyelids, nose or lips. If the tumor involves these structures, portions of them may have to be removed with resulting cosmetic or functional deformities. Furthermore, repairing the resulting defect may involve some of these structures.
- **Rarely**, wounds become infected (fewer than 1%) and require antibiotic treatment. If you are at particular risk for infection, you may be given an antibiotic during surgery.
- There may be excessive bleeding from the wound. Such bleeding can usually be controlled during surgery. There may also be bleeding after surgery. We have never had a significant amount of blood loss, but bleeding into a sutured graft or flap may inhibit good wound healing.
- There may be an adverse reaction to medications used. We will carefully screen you for any history of problems with medications; however, new reactions to medications may occur.
- There is a small chance that your tumor may regrow after surgery. Previously treated tumors and large, longstanding tumors have the greatest chance for recurrence.

Locations



The Bethesda surgery center is located in the Champlain Building at the corner of Democracy Boulevard and Rockledge Drive, near Montgomery Mall.



The Virginia surgery center is located just west of I-495 at the corner of Arlington Blvd and Prosperity Ave (Exit 50A from the beltway).

Dr. Mark Welch was graduated from the New York Medical College and served for thirteen years in the U.S. Army Medical Corps. He completed a residency in Dermatology at Walter Reed Army Medical Center and a fellowship in Mohs Micrographic Surgery at Brooke Army Medical Center.

Dr. Andrew Montemarano is a graduate with honors from the Philadelphia College of Osteopathic Medicine. He served 9 years in the U.S. Army Medical Corps as a Brigade Surgeon in the Republic of South Korea, clinical investigator at the Walter Reed Army Institute of Research, and staff dermatologist at the Walter Reed Army Medical Center. He has completed a fellowship in Mohs Micrographic Surgery under the direction of Donald J. Grande, M.D. of Boston University Medical School.

Dr. Charles Toner is a graduate of the Uniformed Services University of the Health Sciences and retired from the U.S. Navy after 29 years of service. He last served as Chairman of Dermatology and Chief of Mohs Surgery at the National Naval Medical Center in Bethesda and as Dermatology Consultant to the U.S. Congress. He has completed a fellowship in Mohs Micrographic Surgery at the Skin Cancer Center in Winston-Salem, North Carolina.

Dr. Rosemarie Liu received her M.D. degree from Eastern Virginia Medical School. Following an internship at Yale-New Haven Medical Center, she completed a dermatology residency and procedural dermatology (Mohs) fellowship at the University of Wisconsin (UW) Hospital and Clinics in Madison, WI. After serving as co-chief resident, she was a clinical instructor in the UW Department of Dermatology.

Dr. Jason Marquart is a graduate of the Uniformed Services University of the Health Sciences. He completed his dermatology residency at the Walter Reed Army Medical Center and the National Naval Medical Center. He has completed a fellowship in Mohs Micrographic Surgery at the Cooper University Hospital, Robert Wood Johnson Medical School in Marlton, NJ. He is an assistant professor at the Uniformed Services University of the Health Sciences and Howard University Hospital.

All doctors are Board Certified in dermatology and are members of the American Academy of Dermatology and the American College of Mohs Surgery. Only **fellowship-trained** surgeons are admitted to the ACMS.

Together they have over seventy years of medical experience and have treated more than 70,000 skin cancers.

Important Reminders

- **DO** advise us as soon as possible if you must cancel or change your appointment.
- **DO** bring your insurance card(s) and a valid picture identification.
- **DO** plan on being in the office for 3-6 hours.
- **DO** take your usual medications on schedule unless instructed otherwise.
- **DO** eat a big breakfast, unless instructed otherwise.
- **DO** have Tylenol at home for post-surgical pain.
- **DO** let our staff know if you take Coumadin, Plavix or antibiotics before dental work.
- **DO** plan to return 5-7 days after your surgery.
- **DO NOT** plan on exercising for 1 week after surgery.
- **DO NOT** take aspirin, any aspirin-containing products (unless prescribed by a physician), or Vitamin E for ten days prior to the surgery. Please read the label on all over-the-counter medicines.
- **DO NOT** wear fragrances or perfumes.
- **DO NOT** consume alcohol 24 hours prior to or 48 hours after surgery.

Finally...

Please review this handout. We want you to be as comfortable, relaxed and informed as possible.

Mark L. Welch, M.D.
Andrew D. Montemarano, D.O.
Charles B. Toner, M.D.
Rosemarie H. Liu, M.D.
Jason D. Marquart, M.D.

The Champlain Building
 Skin Cancer Surgery Center
 6410 Rockledge Drive, #300
 Bethesda, MD 20817-7811
 301-564-3131 tel
 301-564-6391 fax

Prosperity Medical Center
 Skin Cancer Surgery Center
 8503 Arlington Blvd, #150
 Fairfax, VA 22031-4603
 703-204-0000 tel
 703-560-3960 fax

www.mohssurgerycenter.net

Copyright 2011, all rights reserved. (rev09.14)